

AIROMIR AUTOHALER DEVICE SALBUTAMOL

PACKAGE LEAFLET: INFORMATION FOR THE USER

Read all of this leaflet carefully before you start taking this medicine.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.
- If any of the side effects get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

IN THIS LEAFLET:

- 1. What Airomir Autohaler device is and what it is used for**
- 2. Before you use Airomir Autohaler device**
- 3. How to use Airomir Autohaler device**
- 4. Possible side effects**
- 5. How to store Airomir Autohaler device**
- 6. Further information**

1. WHAT AIROMIR AUTOHALER DEVICE IS AND WHAT IT IS USED FOR

Airomir Autohaler device contains salbutamol, which is a bronchodilator. It opens up the breathing tubes of the lungs and makes breathing easier. Salbutamol can help you feel less tight-chested and less breathless or wheezy.

Airomir Autohaler device is used:

- to treat asthma by relieving wheezing and shortness of breath
- to prevent asthma following exercise
- for the treatment of wheezing and shortness of breath caused by certain other chest diseases.

2. BEFORE YOU USE AIROMIR AUTOHALER DEVICE

DO NOT use Airomir Autohaler device and consult your doctor if you:

- are allergic (hypersensitive) to salbutamol or any of the other ingredients of this medicine

Airomir Autohaler device should NOT be used to treat premature labour or threatened miscarriage, as inhaled salbutamol products are not suitable for this purpose.

Take special care with Airomir Autohaler device

Talk to your doctor before you start to use this medicine if you:

- have thyroid problems
- have severe asthma
- have hypoxia (lack of oxygen in the body)
- have a history of heart disease or angina..

Taking other medicines

Talk to your doctor if you are taking any of the following:

- xanthines e.g. aminophylline or theophylline
- metronidazole
- disulfiram
- steroids
- water tablets (diuretics)
- long-term laxatives
- beta-blockers e.g. propranolol.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including any medicines obtained without a prescription.

Important information about some of the ingredients of Airomir Autohaler device

- Airomir Autohaler device contains a small amount of ethanol (alcohol).

Pregnancy and breast-feeding

- If you are pregnant, planning to become pregnant or are breast-feeding, ask your doctor for advice before taking this medicine.

Driving and using machines

- Airomir Autohaler device may cause dizziness. If you are affected, DO NOT drive or operate machinery.

3. HOW TO USE AIROMIR AUTOHALER DEVICE

Always use Airomir Autohaler device exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

If you see any other doctor or dentist, please tell them that you are using Airomir Autohaler device.

The usual dose is:

Adults (including the elderly)

- **For the relief of symptoms such as wheezing and shortness of breath**
Either one or two puffs as needed.
- **For asthma due to exercise**
Two puffs before exercising.

Children

- **For the relief of symptoms such as wheezing and shortness of breath**
Usually one puff as needed. This may be increased to two puffs if necessary.
- **For asthma due to exercise**
Usually one puff before exercising. This may be increased to two puffs if necessary.

Children using this inhaler should be supervised by an adult and should use this inhaler only as advised by the doctor.

All patients

- **If you have taken a dose of two puffs, you should wait four hours before taking another dose. DO NOT take more than eight puffs in 24 hours.**

- **If you are not sure how many puffs to take or how often to take them, please ask your doctor or pharmacist.**
- **If your usual treatment is not working or you need more than eight puffs per day, please tell your doctor. You should NOT increase your dose without consulting your doctor.**

If you use more of your Airomir Autohaler device than you should

If you accidentally take more puffs than your doctor told you to, please contact your nearest hospital casualty department, or tell your doctor, immediately. An overdose is likely to cause agitation and fast heart beat.

If you stop using Airomir Autohaler device

DO NOT stop taking your medicine without talking to your doctor first even if you feel better.

Before use:

This inhaler has a different feel and taste to CFC inhalers you may have used before.

If this is a new device or if you have not used it for 2 weeks or more, it must be tested before use by releasing 4 puffs into the air in the following way:

1. Take the cover off from the mouthpiece, by pulling down on the lip at the back (fig. 1).
2. Shake the inhaler vigorously. Point the mouthpiece away from you so that the puffs of medicine will go into the air. Hold the inhaler upright and push the lever up so that it stays up (fig. 2). 3. Then, to release a puff, push the dose release slide on the bottom of the device in the direction indicated by the arrow on it (fig. 3).
4. To release the next puff you need to return the lever to its down position (fig. 4) and follow steps 2 and 3 again. Steps 2 and 3 should be repeated until a total of 4 puffs have been released. After releasing the fourth puff, return the lever to its down position, to be ready to take a puff of your medicine.

DO NOT USE THE DOSE RELEASE SLIDE TO TAKE YOUR MEDICINE, THE AUTOHALER DEVICE WILL AUTOMATICALLY RELEASE A DOSE WHEN YOU BEGIN TO BREATHE IN FROM THE MOUTHPIECE.

The instructions for taking a puff are given over the page.

How to use your Airomir® Autohaler® device

1. Take the cover off from the mouthpiece, by pulling down on the lip at the back.
2. Shake the inhaler vigorously. Hold your Autohaler device upright as shown. Push the lever up so that it stays up. Continue to hold your Autohaler device upright, making sure that your hand is not blocking the air vent (marked by **X** in fig. 2) at the bottom of the device.
3. Breathe out as far as is comfortable and then immediately place the mouthpiece in your mouth and close your lips around it.
4. Breathe in slowly and deeply through the mouthpiece. Do not stop breathing in when you hear the slight click and feel the puff in your mouth as it is important that you carry on breathing in after the puff is released.
5. Hold your breath for 10 seconds and then breathe out slowly. 6. The lever must be lowered to the down position immediately after each puff whilst holding the Autohaler device upright. If your doctor has prescribed more than one puff, repeat steps 2 - 6. After use, replace the cover on the mouthpiece.

Cleaning instructions

For normal hygiene, the mouthpiece of your Autohaler device should be cleaned weekly with a clean, dry tissue or cloth.

CAUTION: Do not push a drying cloth or anything else into any part of the device since it may cause damage to its operating parts.
Do not take the device apart.

How to tell when your device is empty

When the device is completely empty you will not feel or hear any propellant being discharged.

Further questions

If you have any further questions on the use of this product, ask your doctor or pharmacist.

4. POSSIBLE SIDE EFFECTS

Like all medicines, Airomir Autohaler device can cause side effects, although not everybody gets them.

If, immediately after inhaling a puff, your asthma worsens, do not use Airomir Autohaler device again and contact your doctor straight away.

If the following happens, stop using Airomir Autohaler device and tell your doctor immediately or go to the casualty department at your nearest hospital:

- an allergic reaction (swelling of the lips, face or neck leading to severe difficulty in breathing; skin rash or hives).

This is a very serious but rare side effect. You may need urgent medical attention or hospitalisation.

The following side effects have been reported:

- feeling sick
- being sick
- a dry, sore mouth
- dizziness
- headache
- mild tremor (shaking)
- muscle cramps
- tachycardia (fast heart rate) or palpitations
- hyperactivity may occur rarely in children.

Using Airomir Autohaler may lead to low potassium levels in your blood. Your doctor may wish you to have regular blood tests to check your blood potassium levels.

Although it is not known exactly how often this happens, some people may experience chest pain (due to heart problems such as angina). Tell your doctor as soon as possible if you develop these symptoms whilst receiving treatment with Airomir, but do not stop taking this medicine unless told to do so.

If any of the side effects continue for more than a few days, get serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

5. HOW TO STORE AIROMIR AUTOHALER DEVICE

Keep out of the reach and sight of children.

Store this inhaler below 30°C. Avoid storage in direct sunlight or heat. Protect from frost.

The metal canister is pressurised. Do not attempt to puncture it or burn it, even when empty.

Do not use Airomir Autohaler device after the expiry date that is stated on the outer packaging. The expiry date refers to the last day of that month.

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

6. FURTHER INFORMATION

Airomir Autohaler device does not contain CFC's; instead, the inhaler contains a hydrofluoroalkane (HFA-134a) as an inactive ingredient. Chlorofluorocarbons (CFC's) have been shown to damage the ozone layer in the atmosphere. HFA's have been developed as a replacement for CFC's because they do not damage the ozone layer.

What Airomir Autohaler device contains:

- The active ingredient is salbutamol.
Each puff contains salbutamol sulphate equivalent to salbutamol 100 micrograms.
- The other ingredients are oleic acid, ethanol, propellant HFA-134a. Propellant HFA-134a is a new propellant which has been developed to replace chlorofluorocarbons (CFCs). This inhaler does not contain CFC propellants.

What Airomir Autohaler device looks like and contents of the pack:

- The Airomir Autohaler device is a breath actuated pressurised inhalation suspension.
- The inhaler contains 200 puffs.

Marketing Authorisation Holder and Manufacturer

The marketing authorisation holder is Teva UK Limited, Brampton Road, Hampden Park, Eastbourne, East Sussex, BN22 9AG, United Kingdom.

Airomir Autohaler devices are made by 3M Health Care Limited, Loughborough, Leicestershire LE11 1EP, England.

This leaflet was last revised: December 2009

PL 00289/1411